

Report Documentation Page				Form Approved OMB No. 0704-0188	
Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.					
1. REPORT DATE <b>01 JAN 2011</b>		2. REPORT TYPE <b>N/A</b>		3. DATES COVERED <b>-</b>	
4. TITLE AND SUBTITLE <b>Real-time Calculation of System-level Complexity During Trauma/Hemorrhage: Can We Do It?</b>				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S) <b>Batchinsky A. I., Baker W. L., Isbell C. L., Necsoiu C., Walker 3rd K. P., Marczyk J., Salinas J., Cancio L. C.,</b>				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) <b>United States Army Institute of Surgical Research, JBSA Fort Sam Houston, TX</b>				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT <b>Approved for public release, distribution unlimited</b>					
13. SUPPLEMENTARY NOTES					
14. ABSTRACT					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT <b>UU</b>	18. NUMBER OF PAGES <b>1</b>	19a. NAME OF RESPONSIBLE PERSON
a REPORT <b>unclassified</b>	b ABSTRACT <b>unclassified</b>	c THIS PAGE <b>unclassified</b>			

**"Real-time calculation of system-level complexity during trauma/hemorrhage: can we do it?"**

**Andriy Batchinsky<sup>1</sup>**, William Baker<sup>1</sup>, Claire Isbell<sup>1</sup>, Corina Necsoiu<sup>1</sup>, Kerfoot Walker<sup>1</sup>, Jacek Marczyk<sup>2</sup>, José Salinas<sup>1</sup>, Leopoldo Cancio<sup>1</sup>. <sup>1</sup> US Army Inst of Surgical Research, Fort Sam Houston, TX; <sup>2</sup> Ontonix S.r.l., Como, Italy.

**Objectives:** We previously showed that sample entropy (SampEn), and other nonlinear measures of the complexity of the ECG time series, decrease in response to hypovolemia and/or injury. These measures characterize only one signal, and thus refer to *signal-level complexity*. In contrast, *system-level complexity* quantifies the amount of interaction among the signals arising from all available sensors attached to a patient (ECG, blood pressure, oxygen saturation, etc.). We found that system-level complexity (OSC) calculated with OntoSpace software (Ontonix S.r.l., Como, Italy) fluctuates during critical events such as asphyxia. Relatedly, *system robustness* (OSR) represents the margin between current OSC and maximal or minimal possible OSC. OSR thus is greatest when OSC is in the middle range; it decreases when OSC approaches either extreme, where the system becomes unstable and is prone to crash. Here, we present data calculated in real time from both signal-level and system-level complexity in a model of acute respiratory distress syndrome (ARDS) due to trauma, hemorrhagic shock and resuscitation. **Methods:** Nine swine were anesthetized with ketamine and midazolam and underwent baseline measurements (BL), right-chest pulmonary contusion (PC), hemorrhage of 12 mL/kg (Bleed), resuscitation with lactated Ringer's (LR), transfusion of shed blood (Tx), and post-resuscitation observation (Post-Resus). Data were collected continuously and analyzed in 15-min datasets. We calculated heart rate (HR, bpm); mean arterial pressure (MAP, mmHg); PaO<sub>2</sub>-to-FiO<sub>2</sub> ratio (PFR); ECG SampEn (unitless); ECG multiscale entropy (MSE, unitless), and percentage of normal-to-normal RRs differing by more than 50 ms (pNN<sub>50</sub>). We calculated OSC and OSR (both unitless) from 56 different channels of single-sensor data. **Results:** see table. Means±SEMs are reported. Statistics: one way ANOVA with Tukey's adjustment.

	BL	PC	Bleed	LR	Tx	Post Resus
HR	80±5	101±5*	109±6*	115±12*	106±9*	104±7*
MAP	83±5	54±4*	58±6*	68±4*†	69±4*†	75±4†
PFR	479±9	198±29*	196±36*	162±36*	174±35*	246±39*¥#°
SampEn	2.0±0.2	1.6±0.2*	1.5±0.1*	1.6±0.2	1.7±0.3	2.0±0.2¥
MSE	18.9±2.4	15.3±2.4	13.5±1.8	14.1±2.3	14.1±2.6*	16.8±2.4
PNN50	0.28±0.10	0.10±0.04	0.09±0.03	0.11±0.02	0.15±0.05	0.22±0.06†¥
OSC	3.6±0.5	20.0±1.9*	12.7±1.5*†	9.2±0.9*†¥	6.5±0.6*†¥	9.5±0.9*†¥#
OSR	93.5±0.3	61.1±1.3*	78.5±0.9*†	82.9±0.6*†	88.7±0.32†°	83.2±0.68*†#

\*Significant (p<0.05) vs.BL; †vs. PC; ¥vs. Bleed; °vs. LR; #vs. Tx.

**Conclusions:** Measures of signal-level complexity like SampEn, and measures of system-level complexity like OSC, address fundamentally different characteristics of a patient's physiology. This is the first report containing data acquired in real-time and combining both approaches in a model of ARDS caused by trauma/hemorrhage. Future work will address whether this combined approach to monitoring can be used to improve outcomes in critical care by enabling earlier or more effective intervention in potentially unstable patients.